

(OLD COURSE)

QP Code : MV-17726

(3 Hours)

[Total Marks : 100

- N. B. : (1) Question No. 1 is compulsory.
 (2) Attempt any four questions from the remaining six questions.
 (3) Figures to the right hand side indicate full marks.

1. (a) Solve $(2D^2 + 5D - 12)y = 0$ 3
 (b) Evaluate $\int_0^{\infty} x e^{-x^4} dx$ 3
 (c) Evaluate $\int_0^1 \int_{x^2}^x xy(x+y) dy dx$ 3
 (d) Evaluate $\int_0^1 \sqrt{1-x^4} dx$ 3
 (e) If $B(n,3) = \frac{1}{105}$ find 'n' where n is +ve integer. 4
 (f) Using Euler's method find approximate value of y at x=1 in five steps taking 4

n = 0.2. Given $\frac{dy}{dx} = x+y$ and $y(0) = 1$.

2. (a) Prove that $\int_0^3 \frac{x^{\frac{3}{2}}}{\sqrt{3-x}} dx \cdot \int_0^1 \frac{dx}{\sqrt{1-x^4}} = \frac{432}{35} \pi$ 6
 (b) Solve $(D^2 - D + 1)Y = \cos 2x$. 6
 (c) Solve the differential equation $\frac{dy}{dx} = xy$ with initial conditions $y(1) = 2$ and 8
 find y at (i) $x = 1.2$ (ii) $x = 1.4$. By Runge-Kutta method of Fourth order.

3. (a) In a circuit containing inductance, L, resistance R, voltage E, the current I, is 6
 given by $L \frac{dI}{dt} + RI = E$, find current I at time t if at $t = 0$, $I = 0$ and L, R, E are
 constant.
 (b) Find the length of the astroid $x = a \cos^3 t$, $y = a \sin^3 t$ 6
 (c) Solve $\frac{d^2y}{dx^2} - 4 \frac{dy}{dx} + 4y = x^2 + e^x + \cos^2 x$ 8

[TURN OVER

Con. 9418-14.



4. (a) Evaluate $\int_0^a \int_0^a \int_0^a (yz + zx + xy) dx dy dz$ 6

(b) Find the volume bounded by the cone $z^2 = x^2 + y^2$ and the paraboloid; 6

$$z = x^2 + y^2$$

(c) Apply the method of variation of parameters to solve 8

$$(D^2 - 2D + 2)y = e^x \tan x.$$

5. (a) Solve using Taylor's series method the differential equation $\frac{dy}{dx} = x + y$ start 6

from $x = 1, y = 0$ and carry to $x = 1.2$ with $h = 0.1$

(b) Find by double integration the area of the cardioid $r = a(1 + \cos \theta)$ 6

(c) Change the order of integration and evaluate 8

$$\int_0^a \int_{\frac{x}{a}}^{2a-x} xy \, dy \, dx$$

6. (a) Change to polar co-ordinates and evaluate 6

$$\int_0^a \int_y^a \frac{x^2}{\sqrt{x^2 \times y^2}} dx dy$$

(b) Find the length of the cardioid $r = a(1 - \cos \theta)$ which lies out side the circle 6

$$r = a \cos \theta.$$

(c) Prove that $B(m, n) = \frac{\Gamma(m) \Gamma(n)}{\Gamma(m+n)}$ 8

7. (a) Solve $(x^2 + y^2 + 1) dx - 2xy dy = 0$ 6

(b) Solve $\frac{dy}{dx} = x^3 y^3 - xy$ 6

(c) Solve $(D^2 + 2)y = e^x \cos x + x^2 e^{3x}$ 8

